



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

## Clinical Communications

### COVID-19 in a series of patients with aspirin-exacerbated respiratory disease

Andrew A. White, MD<sup>a</sup>, Katherine N. Cahill, MD<sup>b</sup>,  
Elina Jerschow, MD<sup>c</sup>, Merin Kuruvilla, MD<sup>d</sup>,  
Esha Sehanobish, PhD<sup>c</sup>, Jillian Bensko, PA-C<sup>e</sup>,  
Tanya M. Laidlaw, MD<sup>e</sup>, and Joshua M. Levy, MD, MPH<sup>f</sup>

#### Clinical Implications

- We report a series of patients with aspirin-exacerbated respiratory disease who became infected with severe acute respiratory syndrome coronavirus 2. Most of these patients had been treated with biologics for their baseline respiratory disease and did not appear to experience increased morbidity related to coronavirus 2019.

Early in the severe acute respiratory syndrome coronavirus (SARS-CoV-2) pandemic, asthma was understood to be a risk factor for poor outcomes in those infected. Yet, as understanding of the disease matured, asthma was not consistently observed as a risk factor. Because asthma is a heterogeneous disorder, asthma endotypes may impact coronavirus 2019 (COVID-19)-related morbidity.

Aspirin-exacerbated respiratory disease (AERD) is an inflammatory disorder characterized by nasal polyposis and asthma, defined by hypersensitivity to nonsteroidal anti-inflammatory drugs, and is associated with high levels of type 2 inflammation involving mast cell and eosinophil activation. AERD is enriched in populations with severe asthma and is also recognized as the most likely endotype to experience severe and early nasal polyp recurrence, making this group of patients much more difficult to control.<sup>1</sup> Given this burden of illness, we evaluated the morbidity associated with SARS-CoV-2 infection in our respective AERD cohorts.

A total of 19 patients with AERD were identified with SARS-CoV-2 infection documented by PCR, convalescent antibody testing, or compatible clinical syndrome after close exposure to a known case (Table 1). Respiratory biologic use was common (15 of 19) and was continued throughout infection with SARS-CoV-2. All but 1 patient met criteria for severe asthma based on FEV<sub>1</sub> impairment or need for biologic or systemic corticosteroid therapy. Of the 16 symptomatic infections, 2 patients required hospitalization, and all other patients experienced mild to moderate self-limited symptoms. Some reported residual health effects between 2 and 11 months from infection. Our cohort sample is not large enough to analyze statistically, but a few cases merit closer review.

A 53-year-old obese man was on baseline dupilumab therapy when he developed severe COVID-19 pneumonia and a moderate asthma exacerbation. He was hospitalized for severe hypoxia complicated by a pulmonary embolism. Before hospitalization he had impairment in FEV<sub>1</sub> (77%), a normal Sino-Nasal Outcome Test-22 score of 7, and an elevated

absolute eosinophil count of 980 cells/ $\mu$ L. While hospitalized, he was treated with remdesivir, dexamethasone, supplemental O<sub>2</sub>, and enoxaparin sodium and then was transitioned to oral apixaban for the pulmonary embolism. At the time of publication, he has returned to his pre-COVID baseline medications and oral apixaban.

A 75-year-old woman with diabetes on anticoagulation was hospitalized. At baseline, she was treated with benralizumab and had received 2 systemic corticosteroid bursts in the year before infection. Baseline FEV<sub>1</sub>, asthma control test score of 20, and Sino-Nasal Outcome Test-22 score suggested reasonable baseline airway control. She was treated with remdesivir and dexamethasone and at the time of publication is still requiring home oxygen.

A 54-year-old diabetic woman with hypertension, morbid obesity, corticosteroid-dependent asthma, impaired baseline lung function on dupilumab, and a long-term taper of hydrocortisone contracted SARS-CoV-2 and was treated with a prednisone taper and doxycycline for COVID-19 pneumonia. She recovered after 2 weeks and did not require supplemental oxygen or hospitalization.

A 55-year-old man with hypereosinophilia and coronary artery disease was on maintenance treatment with benralizumab. He had required 6 courses of systemic corticosteroids in the previous year due to poor airway control and asthma exacerbations. His COVID-19 infection was asymptomatic.

A 74-year-old man with Parkinson disease, coronary artery disease, and baseline impairment in lung function was on dupilumab at the time of his COVID-19 infection. He experienced cough and worsening shortness of breath as well as significant fatigue. Despite this, he did not require oxygen or hospitalization, nor did he develop any severe complication.

Type 2 inflammation may influence the course of COVID-19 in several ways. SARS-CoV-2 uses the angiotensin-converting enzyme 2 (ACE2) receptor for cell entry, a receptor present throughout the respiratory epithelium, with host transmembrane protease serine 2 (TMPRSS2) providing cleavage of the spike protein, which is required for virus fusion. These proteins are influenced by type 2 immune processes, with IL-13 decreasing ACE2 levels and increasing TMPRSS2 levels.<sup>2</sup> Consistent with the hypothesis that high type 2 inflammation could have a modifying effect, patients with asthma presenting to the emergency department with COVID-19 and a prior absolute eosinophil count of 150 cells/ $\mu$ L were less likely (odds ratio, 0.46) to be admitted to the hospital. Hospitalized patients with absolute eosinophil count of less than 150 cells/ $\mu$ L were also significantly less likely to die (odds ratio, 0.006).<sup>3</sup> Most of our cohort members were on an asthma biologic (79%) yet experienced predominantly mild infection, suggesting that the underlying disease process might be protective independent of biologic-induced dampening of the type 2 signal. Yet, in another study, there did not appear to be a difference in ACE2 and TMPRSS2 expression in patients with asthma versus healthy controls, but male sex, Black/African American race, and diabetes mellitus were associated with higher ACE2 and TMPRSS2 gene expression.<sup>4</sup>

**TABLE I.** Clinical characteristics of SARS-CoV-2 infection in AERD

Age/sex	Ethnicity	Comorbidities*	Biologics/systemic corticosteroids/inhaler type/LTMD	Aspirin dose <sup>†</sup> /duration	Sinus surgery/recent surgery <sup>‡</sup>	SNOT-22 score	FEV <sub>1</sub> (% pred)	AEC (cells/ $\mu$ L)	COVID outcome	"Long-haul" symptoms
17F	Non-Hispanic White	None	ICS	1300mg 10 mo	1 sx Not recent	6	98	730	Mild outpatient	N
18F	Non-Hispanic White	None	Dupilumab, ICS/LABA	N	1 sx	47	91	170	Mild outpatient	N
36F	Non-Hispanic White	Hypothyroid	Dupilumab Methylprednisolone, ICS/ LABA, LTMD	N	0	52	81	NA	Moderate emergency room	N
36M	Black/AA	HTN, obesity	Dupilumab, hydrocortisone, ICS/ LABA	N	5 sx Not recent	NA	54	1600	Asymptomatic	N
37F	Non-Hispanic White	None	Benralizumab, ICS/ LABA, LTMD	650 mg 3 y	2 sx Not recent	13	97	NA	Mild outpatient	N
37F	Non-Hispanic White	Pre-DM, morbid obesity, OSA	Dupilumab, ICS/LABA	N	2 sx Not recent	60	111	340	Moderate outpatient	Y, Fatigue
39F	Non-Hispanic White	CSU, chronic otitis media, psoriasis	Dupilumab Omaliuzumab No other controllers	325 mg 3 y	7 sx Not recent	31	91	560	Mild outpatient	N
42M	Asian	Hypothyroid	ICS/LTMD	1300mg 9 y	1 sx Not recent	64	52	460	Mild outpatient	N
43M	Non-Hispanic White	Non	Dupilumab, ICS/LABA, LTMD	1300mg 2 y	2 sx Not recent	8	92	650	Mild outpatient	Y, Fatigue
50F	Non-Hispanic White	HTN, pre-DM	ICS/LABA	N	1 sx Recent	52	93	280	Mild outpatient	Y, Fatigue
50F	Non-Hispanic White	None	Mepolizumab, ICS/ LABA, LTMD	650 mg 2 y	0 surgery	10	97	NA	Mild outpatient	N
50F	Non-Hispanic White	HTN	ICS/LABA	1300mg 10 y	0 surgery	NA	57	1350	Moderate outpatient	N
53F	Non-Hispanic White	None	Dupilumab, ICS/LABA	325 mg 3 y	4 sx Not recent	NA	52	900	Asymptomatic	N
53M	Non-Hispanic White	Obesity	Dupilumab, ICS/LABA	N	2 sx Not recent	7	77	980	Severe, hospitalized, COVID- related pneumonia, pulmonary embolus	Y, Fatigue
54F	Black/AA	DM, obesity	Dupilumab, hydrocortisone, ICS/ LABA	N	10 sx > Not recent	NA	75	1500	Moderate Outpatient	Y, Fatigue
55M	Non-Hispanic White	HTN, CAD, hypereosinophilia, food allergy	Benralizumab ICS/LABA/LTMD	1300mg 11 mo	1 sx Not recent	1	80	0	Asymptomatic	N
57F	Black/AA	None	Dupilumab, ICS/LABA	N	3 sx Not recent	NA	74	200	Moderate outpatient	N

74M	Non-Hispanic White	Parkinson, CAD, HL	Dupilumab, ICS/LABA/LAMA	325 mg >10 y	1 sx Not recent	NA	58	NA	Moderate outpatient	N
75F	Non-Hispanic White	HTN, DM, HL, Afib	Benralizumab, ICS/LABA, L/TMD	N	2 sx Not recent	9	102	0	Severe hospitalized	Y, Oxygen dependent

AA, African American; AEC, absolute eosinophil count; Afib, atrial fibrillation; CAD, coronary artery disease; CSU, chronic spontaneous urticaria; DM, diabetes mellitus; HL, hyperlipidemia; HTN, hypertension; ICS, inhaled corticosteroid; LABA, long-acting beta-agonist; L/TMD, leukotriene modifier drug; N, no; *pre-DM*, pre-diabetes mellitus; *pred*, predicated; OSA, obstructive sleep apnea; *SNOT-22*, Sino-Nasal Outcome Test 22; Y, yes.

Mild = significant COVID-19 symptoms < 7 d, no need for significant medical intervention.

Moderate = significant COVID-19 symptoms ≥ 7 d and/or need for emergency room care or multiple outpatient medical visits.

Severe = hospitalization/intensive care, and/or oxygen requirement.

\*All patients had a confirmed diagnosis of AERD including sinonasal polyposis, persistent asthma, and nonsteroidal anti-inflammatory drug reactivity. Data reported represent the last reported value before COVID-19 diagnosis.

†Aspirin dose is typically divided in half and taken twice daily, except 325 mg, which is given once daily.

‡Recent surgery = sinus surgery within 6 mo of COVID-19 infection.

Asthma exacerbations are frequent after viral respiratory infections. Whether this would be true following SARS-CoV-2 infections was the topic of much discussion, yet current data suggest that asthma exacerbations from SARS-CoV-2 are not a common contributor to morbidity.<sup>5</sup> Notably, our cohort with moderate to severe asthma required few systemic corticosteroids following COVID-19 illness.

Asthma is not overrepresented in cohorts of patients with severe SARS-CoV-2 infection.<sup>6</sup> Yet, some data do generate concern regarding the outcomes in asthmatic and allergic patients. In a Korean cohort study of 219,959 subjects who underwent COVID-19 testing, asthma and allergic rhinitis were associated with increased risk of test positivity and worsened clinical outcomes.<sup>7</sup> A similarly large database (>17 million patients) noted an increased risk of COVID-19–related death in patients with asthma, but only those with recent oral corticosteroid use.<sup>8</sup> In a review of 634 Dutch patients with asthma on respiratory biologic therapy during the initial wave of the pandemic, 9 were diagnosed with COVID-19 of which 7 required hospitalization, 5 required intubation, and 1 patient died. These reports suggest that COVID-19 morbidity in patients with severe asthma on biologic therapy could be quite high.<sup>9</sup> Regional and temporal differences in testing availability for asymptomatic or mild cases, treatment and health care resource availability, and prevalence of high-risk comorbidities make generalizability of these reports difficult.

Collectively, the authors follow a large cohort of approximately 1000 patients with AERD. In these combined cohorts, COVID-19 infection did not appear to have higher morbidity than what is seen in the general population. This is reassuring in a group of patients with exacerbation-prone asthma. It is intriguing that a strong type 2 signal might be protective—a signal that the vast majority of patients with AERD exhibit. Yet, our cohort is not large enough to draw firm conclusions. Whether specific treatments that skew type 2 immune patterns might be influential in SARS-CoV-2 infection remains to be seen.

<sup>a</sup>Division of Allergy, Asthma and Immunology, Scripps Clinic, San Diego, Calif

<sup>b</sup>Division of Allergy, Pulmonary, and Critical Care Medicine, Vanderbilt University Medical Center, Nashville, Tenn

<sup>c</sup>Department of Medicine, Allergy/Immunology Division, Montefiore Medical Center/Albert Einstein College of Medicine, Bronx, NY

<sup>d</sup>Division of Pulmonary, Allergy, Critical Care, and Sleep Medicine, Emory University, Atlanta, Ga

<sup>e</sup>Department of Medicine, Harvard Medical School, the Division of Allergy and Clinical Immunology, Brigham and Women's Hospital, Boston, Mass

<sup>f</sup>Department of Otolaryngology – Head and Neck Surgery, Emory University, Atlanta, Ga

No funding was received for this study.

Conflicts of interest: A. A. White is on speakers bureau for AstraZeneca, Regeneron/Sanofi, and Optinose; on Advisory Board for Genentech, Regeneron/Sanofi, Optinose, and GlaxoSmithKline; and has received grant support from AstraZeneca. K. N. Cahill is on Advisory Board for Teva, Novartis, GlaxoSmithKline, Blueprint Medicines, Regeneron, and Sanofi-Pasteur; and has received consulting fees from Third Harmonic Bio, Novartis, and Ribon Therapeutics. E. Jerschow is on Advisory Board for GlaxoSmithKline, Sanofi-Regeneron, and Novartis/Genentech; and has received grant support from AstraZeneca and Cumberland. T. M. Laidlaw is on Advisory Board for GlaxoSmithKline, Sanofi-Genzyme, and Regeneron; and has received grant support from GlaxoSmithKline and Regeneron. J. M. Levy is on Advisory Board for Regeneron/Sanofi and GlaxoSmithKline; and has received grant support from AstraZeneca, Cumberland Pharmaceuticals, Genentech, Regeneron/Sanofi, and Optinose.

Received for publication February 25, 2021; revised April 24, 2021; accepted for publication April 26, 2021.

Available online ■ ■

Corresponding author: Andrew A. White, MD, Division of Allergy, Asthma and Immunology, Scripps Clinic, 3811 Valley Centre Dr, San Diego, CA 92130.  
E-mail: [White.andrew@scrippshealth.org](mailto:White.andrew@scrippshealth.org).

2213-2198

© 2021 American Academy of Allergy, Asthma & Immunology

<https://doi.org/10.1016/j.jaip.2021.04.060>

#### REFERENCES

1. Levy JM, Rudmik L, Peters AT, Wise SK, Rotenberg BW, Smith TL. Contemporary management of chronic rhinosinusitis with nasal polyposis in aspirin-exacerbated respiratory disease: an evidence-based review with recommendations. *Int Forum Allergy Rhinol* 2016;6:1273-83.
2. Kimura H, Francisco D, Conway M, Martinez FD, Vercelli D, Polverino F, et al. Type 2 inflammation modulates ACE2 and TMPRSS2 in airway epithelial cells. *J Allergy Clin Immunol* 2020;146:80-8.e8.
3. Ferastraoraru D, Hudes G, Jerschow E, Jariwala S, Karagic M, de Vos G, et al. Eosinophilia in asthma patients is protective against severe COVID-19 illness. *J Allergy Clin Immunol Pract* 2021;9:1152-62.e3.
4. Peters MC, Sajuthi S, Deford P, Christenson S, Rios CL, Montgomery MT, et al. COVID-19-related genes in sputum cells in asthma. Relationship to demographic features and corticosteroids. *Am J Respir Crit Care Med* 2020; 202:83-90.
5. Grandbastien M, Piotin A, Godet J, Abessolo-Amougou I, Ederlé C, Enache I, et al. SARS-CoV-2 pneumonia in hospitalized asthmatic patients did not induce severe exacerbation. *J Allergy Clin Immunol Pract* 2020;8: 2600-7.
6. Wang L, Foer D, Bates DW, Boyce JA, Zhou L. Risk factors for hospitalization, intensive care, and mortality among patients with asthma and COVID-19. *J Allergy Clin Immunol* 2020;146:808-12.
7. Yang JM, Koh HY, Moon SY, Yoo IK, Ha EK, You S, et al. Allergic disorders and susceptibility to and severity of COVID-19: a nationwide cohort study. *J Allergy Clin Immunol* 2020;146:790-8.
8. Williamson EJ, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 2020;584:430-6.
9. Eger K, Hashimoto S, Braunstahl GJ, Brinke AT, Patberg KW, Beukert A, et al. Poor outcome of SARS-CoV-2 infection in patients with severe asthma on biologic therapy. *Respir Med* 2020;177: 106287.